



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

D and its extremity B be joined with the points A and C then the totality of triangles will be formed and they will be uniformly distributed on the semicircumference whose diameter is the hypotenuse.

The question is not whether the triangles are uniformly distributed or not but what method gives the *totality* of the series.

Drury College, January 27, 1896.

NOTES.

ERRATA. Professor Beman calls my attention to a manifest error in Professor Klein's paper which I translated for the December number. Vol. II, page 350, should give the series $\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$. The series $1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$

$= \log_e 2$ instead of $\frac{\pi}{4}$.

D. E. SMITH.

Dr. E. A. Bowser writes: Should not problem 43 [Calculus] read $\int_0^1 \frac{x^{a-1} - x^{-a}}{1+x} \frac{dx}{\log x}$, as in Price's Calculus, Vol. II, page 120?

NOTE ON THE SOLUTIONS OF PROBLEM 45, PAGES 274-75.

BY ARTEMAS MARTIN, LL. D.

There is but *one* case in Problem 45, Geometry, *as proposed*. Only the circumscribing circle is required.

The final result may be expressed in the more simple form

$$R = \frac{abc}{2\sqrt{[abc(a+b+c)] - (ab+ac+bc)}}.$$

In the second solution, page 275, the equation

$$\text{"cos } BCA = \cos(BCA + BCO)\text{"}$$

should be

$$\cos BCA = \cos(ACO + BCO).$$